

67 GTG TCA ACC TGG TTG ATA CGT AGT GGT GAA CCT GTG CAA CAC CGC ACT GAA TTC CCG TTC
127 ATC GCA TTC TTA AGC GAG AGA ACA GTT TCA CTA GTT TCA ACG AGA GCA
187 GTA CTC TCA GCT AGT CAT TGT GTT TGC TCA CCA GTG ATT **GGG gtgagatgcgctgtg**
250 aaacacattgtgttgcctaagttaacgcattgttttcacagcagcagtggtttgcgtgcgtgcgtttgtgttctgtatgcgc
329 gcgggtttgcatagaattttaaagtcctatatactttggtcttatattcag GTT TCA TTT CTC ACA CTG AGC AAT
400 GGC GAC CAA CAA GGC ATC CAT CAC CAA CCG TCT GGA GTT AAG GTG GCA CCA GGA TAC ATG
460 CCC TCT TGT ATT TCG GCA CCA CAG AAG AGA CCA ATC CAG ACA CTC AGT GGA TTC GAT
520 ATT GCA TGT GTA ATG CTG GCT CAA ATG GTC AAC TTA CAG AGT GGA ATC AGA GTG ATC AGT
580 CTG CCA CAG CCA TCG GAT ATC CCG CCA CCT GGA ACT GGT GTT TTC ATT GTT GGT TAT GGA
640 AGG GAT GAT AAC GAC GGT GAT CCG TCA CGT AAG AAT GGT GGA ATG AAG **AAA gtgagat**
701 gtgtgtgaaataaacgcacatgactcagtcagtcagatcagatcagtcagtcattctgtgtctgtgtatctgtttgtgt
780 gtctctgtctctacctgactccggtgtgtgtatttggtcagagccttgaataaacacactgtgtttggatgactttgtgtgc
859 agtctcagtagcagagtgatttccatctcggtcattgtgttggtgaggtgaggtgacgtgatgtgaaggtgagttgagga
938 aatggatgggatggatgtgatgtgatgggatgatgatgaccacttgaggagaggaagactcatgaatattctatgcga
1017 gacgatgggaagtgctgtgtgtcatgaagtggggggtcaatgtgttgagattgtgttgaggaggtgtgagatggagah
1096 gtgacttgactcgtcgatatagtgcacatgtgattgtatgtggactattgttgtgtgggtcagtgtgaaggggtggatatt
1175 gtgcacagttgatattctcgaattccactgtgtgtgtttgttttgttcctgtgtgttcttcttccatctgtgactgt
1254 acgctgtgttcagtcactgttctgttggtgtgtgtgtgtgcacacaga GGT CGA GGC ACT ATA ATG CAA TGC
1325 CCA CAT GCG ACC AAT GGC AAT CCT ATA TGT GTG AAA GCA GGT CAG AAT TTC GGA CAG TTA
1385 CCC GCT CCA GGT GAC AGT GGT GGA CCT CTC CTC CCA TCC CTT CAA GGT CCA GTA CTC GGT
1445 GTG GTA TCA CAT GGT GTC ACA CTG CCA AAC CTT CCC GAT ATC ATT GTC GAG TAT GCC AGT
1505 GTC GCT AGA ATG TTG GAT TTT GTA CGC TCC AAT ATT TGA

Fig. 1. Consensus sequence of 3 PCR clones of *Schistosoma mansoni* cercarial elastase gene. Nucleotide numbers correspond to the open reading frame of the cDNA sequence reported by Newport *et al.* (1988). The 2 introns are shown in lower case text. Splice signal and branch formation sequences are shown in underlined text.

[illegible]

Figure 2 - Production of Construct pGEXCEL





Figure 3 - SDS-PAGE Showing Purification of the Recombinant Protein Sm30-GST



M = Protein Mid-Range Molecular Weight Markers

1 = Total proteins from uninduced *E. coli* carrying plasmid pGEXCEL

2 = Total proteins from IPTG-induced *E. coli* carrying pGEXCEL, grown at 30°C

3 = Soluble lysate of IPTG-induced *E. coli* as in lane 2


4 = Fraction containing Sm30-GST following ion-exchange chromatography

5 = Fraction containing Sm30-GST following gel filtration

The recombinant protein Sm30-GST is indicated by "←"

Figure 4:

Western immunoblot showing antibody reactivity of serum from rabbit immunized with Sm30-GST.




Antigens in lanes:

1. Horse albumin
2. Recombinant fusion protein Sm30-GST
3. " " " Sm30Ex1-GST
4. " " " Sm30Ex2-GST
5. " " " Sm30Ex3-GST
6. Recombinant Sj26GST
7. *S. mansoni* cercarial transformation fluid (CTF)

Figure 5


Reactivity of sera from immunized mice against *S. mansoni* cercarial antigens.



A = sera from 6 mice immunized with *S. mansoni* CTF
 B = sera from 6 mice immunized with Sm30-GST
 1 = serum from rabbit BR67
 2 = normal rabbit serum
 3 = larval antigens stained with protogold
 M = molecular weight standards

Figure 6


Reactivity of sera from immunized mice against *S. haematobium* cercarial antigens.



A = sera from 6 mice immunized with *S. mansoni* CTF
 B = sera from 6 mice immunized with Sm30-GST
 1 = serum from rabbit BR67
 2 = normal rabbit serum
 3 = larval antigens stained with protogold
 M = molecular weight standards

Figure 7:

Cross-reactivity of antibodies raised against *S. haematobium* 27kDa larval protease against antigens of *S. mansoni* and *S. margrebowiei*.




Antigens in lanes:

1. *S. mansoni* cercarial antigens
 2. *S. haematobium* cercarial antigens
 3. *S. margrebowiei* cercarial antigens
- M = molecular weight standards (kDa)

Rabbit antiserum used to probe antigens in lanes 1-3 was raised against the 27kDa *S. haematobium* larval protease using the same methods as for BR67, the anti-*S. mansoni* 27kDa larval protease antiserum.

Figure 8:

Antibody reactivity of mice immunized with recombinant fusion proteins against *S. mansoni* cercarial transformation fluid (CTF).



Group	Immunogen
A	Alum and PBS alone (control mice).
B	CTF.
C	Recombinant Sj26.
D	Recombinant Sm30-GST.
E	Recombinant Sm30Ex1-GST
F	Recombinant Sm30Ex2-GST
G	Recombinant Sm30Ex3-GST

Control Sera

- E = Rabbit serum BR67 (anti-27kDa *S. mansoni* larval protease)
- G = Rabbit serum 1093X (anti-*S. japonicum* GST)
- N = Normal rabbit serum

M = Molecular weight markers.

M =